

# 2013年 英文誌掲載論文

Role of sirtuins in kidney disease.	Kitada M, Kume S, Koya D.	Curr Opin Nephrol Hypertens.
Usefulness of a novel system for measuring glucose area under the curve while screening for glucose intolerance in outpatients.	Sakamoto K, Kubo F, Yoshiuchi K, Ono A, Sato T, Tomita K, Sakaguchi K, Matsuhisa M, Kaneto H, Maegawa H, Nakajima H, Kashiwagi A, Kosugi K.	J Diabetes Investig.
A cross-sectional association of obesity with coronary calcium among Japanese, Koreans, Japanese Americans, and U.S. whites.	Fujiyoshi A, Sekikawa A, Shin C, Masaki K, David Curb J, Ohkubo T, Miura K, Kadokawa T, Kadokawa S, Kadota A, Edmundowicz D, Shah A, Evans RW, Bertolet M, Choo J, Willcox BJ, Okamura T, Maegawa H, Murata K, Kuller LH, Ueshima H; ERA JUMP (Electron-Beam Tomography, Risk Factor Assessment Among Japanese and U.S. Men in the Post-World War II Birth Cohort) Study Group.	Eur Heart J Cardiovasc Imaging.
Evaluation of a minimally invasive system for measuring glucose area under the curve during oral glucose tolerance tests: usefulness of sweat monitoring for precise measurement.	Sakaguchi K, Hirota Y, Hashimoto N, Ogawa W, Hamaguchi T, Matsuo T, Miyagawa J, Namba M, Sato T, Okada S, Tomita K, Matsuhisa M, Kaneto H, Kosugi K, Maegawa H, Nakajima H, Kashiwagi A.	J Diabetes Sci Technol.
Effect of sleeve gastrectomy on body weight, food intake, glucose tolerance, and metabolic hormone level in two different rat models: Goto-Kakizaki and diet-induced obese rat.	Trung VN, Yamamoto H, Yamaguchi T, Murata S, Akabori H, Ugi S, Maegawa H, Tani T.	J Surg Res.
Replication study for the association of a single-nucleotide polymorphism, rs3746876, within KCNJ15, with susceptibility to type 2 diabetes in a Japanese population.	Fukuda H, Imamura M, Tanaka Y, Iwata M, Hirose H, Kaku K, Maegawa H, Watada H, Tobe K, Kashiwagi A, Kawamori R, Maeda S.	J Hum Genet.
Gene therapy for neuropathic pain by silencing of TNF- $\alpha$ expression with lentiviral vectors targeting the dorsal root ganglion in mice.	Ogawa N, Kawai H, Terashima T, Kojima H, Oka K, Chan L, Maegawa H.	PLoS One
Hyperglycemia induces abnormal gene expression in hematopoietic stem cells and their progeny in diabetic neuropathy.	Katagi M, Terashima T, Okano J, Urabe H, Nakae Y, Ogawa N, Udagawa J, Maegawa H, Matsumura K, Chan L, Kojima H.	FEBS Lett.
Autophagy: emerging therapeutic target for diabetic nephropathy.	Kume S, Yamahara K, Yasuda M, Maegawa H, Koya D.	Semin Nephrol.
The role of autophagy in the pathogenesis of diabetic nephropathy.	Yamahara K, Yasuda M, Kume S, Koya D, Maegawa H, Uzu T.	J Diabetes Res.
4-Hydroxy hexenal derived from dietary n-3 polyunsaturated fatty acids induces anti-oxidative enzyme heme oxygenase-1 in multiple organs.	Nakagawa F, Morino K, Ugi S, Ishikado A, Kondo K, Sato D, Konno S, Nemoto K, Kusunoki C, Sekine O, Sunagawa A, Kawamura M, Inoue N, Nishio Y, Maegawa H.	Biochem Biophys Res Commun.
Glycated Albumin Predicts the Risk of Mortality in Type 2 Diabetic Patients on Hemodialysis: Evaluation of a Target Level for Improving Survival.	Isshiki K, Nishio T, Isono M, Makiishi T, Shikano T, Tomita K, Nishio T, Kanasaki M, Maegawa H, Uzu T.	Ther Apher Dial.
Obesity-mediated autophagy insufficiency exacerbates proteinuria-induced tubulointerstitial lesions.	Yamahara K, Kume S, Koya D, Tanaka Y, Morita Y, Chin-Kanasaki M, Araki H, Isshiki K, Araki S, Haneda M, Matsusaka T, Kashiwagi A, Maegawa H, Uzu T.	J Am Soc Nephrol.
Replication study for the association of 9 East Asian GWAS-derived loci with susceptibility to type 2 diabetes in a Japanese population.	Sakai K, Imamura M, Tanaka Y, Iwata M, Hirose H, Kaku K, Maegawa H, Watada H, Tobe K, Kashiwagi A, Kawamori R, Maeda S.	PLoS One.
Assessing the clinical utility of a genetic risk score constructed using 49 susceptibility alleles for type 2 diabetes in a Japanese population.	Imamura M, Shigemizu D, Tsunoda T, Iwata M, Maegawa H, Watada H, Hirose H, Tanaka Y, Tobe K, Kaku K, Kashiwagi A, Kawamori R, Maeda S.	J Clin Endocrinol Metab.
Genome-wide association study identifies three novel loci for type 2 diabetes.	Hara K, Fujita H, Johnson TA, Yamauchi T, Yasuda K, Horikoshi M, Peng C, Hu C, Ma RC, Imamura M, Iwata M, Tsunoda T, Morizono T, Shojima N, So WY, Leung TF, Kwan P, Zhang R, Wang J, Yu W, Maegawa H, Hirose H; DIAGRAM consortium, Kaku K, Ito C, Watada H, Tanaka Y, Tobe K, Kashiwagi A, Kawamori R, Jia W, Chan JC, Teo YY, Shyong TE, Kamatani N, Kubo M, Maeda S, Kadokawa T.	Hum Mol Genet.
4-Hydroxy hexenal derived from docosahexaenoic acid protects endothelial cells via Nrf2 activation.	Ishikado A, Morino K, Nishio Y, Nakagawa F, Mukose A, Sono Y, Yoshioka N, Kondo K, Sekine O, Yoshizaki T, Ugi S, Uzu T, Kawai H, Makino T, Okamura T, Yamamoto M, Kashiwagi A, Maegawa H.	PLoS One.
Carotid intima-media thickness and plaque in apparently healthy Japanese individuals with an estimated 10-year absolute risk of CAD death according to the Japan Atherosclerosis Society (JAS) guidelines 2012: the Shiga Epidemiological Study of Subclinical Atherosclerosis (SESSA).	Kadota A, Miura K, Okamura T, Fujiyoshi A, Ohkubo T, Kadokawa T, Takashima N, Hisamatsu T, Nakamura Y, Kasagi F, Maegawa H, Kashiwagi A, Ueshima H; SESSA Research Group; NIPPON DATA80/90 Research Group.	J Atheroscler Thromb.
Enhanced Intestinal Motility during Oral Glucose Tolerance Test after Laparoscopic Sleeve Gastrectomy: Preliminary Results Using Cine Magnetic Resonance Imaging.	Trung VN, Yamamoto H, Furukawa A, Yamaguchi T, Murata S, Yoshimura M, Murakami Y, Sato S, Otani H, Ugi S, Morino K, Maegawa H, Tani T.	PLoS One.
Cilostazol attenuates spontaneous microaggregation of platelets in type 2 diabetic patients with insufficient platelet response to aspirin.	Araki S, Matsuno H, Haneda M, Koya D, Kanno Y, Kume S, Isshiki K, Araki H, Ugi S, Kawai H, Kashiwagi A, Uzu T, Maegawa H.	Diabetes Care.
Ezetimibe prevents hepatic steatosis induced by a high-fat but not a high-fructose diet.	Ushio M, Nishio Y, Sekine O, Nagai Y, Maeno Y, Ugi S, Yoshizaki T, Morino K, Kume S, Kashiwagi A, Maegawa H.	Am J Physiol Endocrinol Metab.
Octreotide improves early dumping syndrome potentially through incretins: a case report.	Sato D, Morino K, Ohashi N, Ueda E, Ikeda K, Yamamoto H, Ugi S, Yamamoto H, Araki S, Maegawa H.	Endocr J.
Ethnic difference in liver fat content: a cross-sectional observation among Japanese American in Hawaii, Japanese in Japan, and non-Hispanic whites in United States.	Azuma K, Curb JD, Kadokawa T, Edmundowicz D, Kadokawa S, Masaki KH, El-Saad A, Nishio Y, Seto TB, Okamura T, Cetinel C, Kadota A, Sutton-Tyrrell K, Miura K, Evans RW, Takamiya T, Maegawa H, Miljkovic I, Kuller LH, Ueshima H, Kelley DE, Sekikawa A.	Obes Res Clin Pract.
LOX-1 ligands containing apolipoprotein B and carotid intima-media thickness in middle-aged community-dwelling US Caucasian and Japanese men.	Okamura T, Sekikawa A, Sawamura T, Kadokawa T, Barinas-Mitchell E, Mackey RH, Kadota A, Evans RW, Edmundowicz D, Higashiyama A, Nakamura Y, Abbott RD, Miura K, Fujiyoshi A, Fujita Y, Murakami Y, Miyamoto N, Kakino A, Maegawa H, Murata K, Horie M, Mitsunami K, Kashiwagi A, Kuller LH, Ueshima H; ERA JUMP Study Group.	Atherosclerosis.
Ketogenic essential amino acids replacement diet ameliorated hepatosteatosis with altering autophagy-associated molecules.	Xu L, Kanasaki M, He J, Kitada M, Nagao K, Jinzu H, Noguchi Y, Maegawa H, Kanasaki K, Koya D.	Biochim Biophys Acta.
Altered unfolded protein response is implicated in the age-related exacerbation of proteinuria-induced proximal tubular cell damage.	Takeda N, Kume S, Tanaka Y, Morita Y, Chin-Kanasaki M, Araki H, Isshiki K, Araki S, Haneda M, Koya D, Kashiwagi A, Maegawa H, Uzu T.	Am J Pathol.